

Fax Referral To: 1-800-323-2445 Phone: 1-800-237-2767 Email Referral To: Customer.ServiceFax@CVSHealth.com



			mitting a Referi	ral	
PATIENT INFORMATION	N (Complete or include d	lemographic sh	ieet)		
Patient Name:			DOB:		_Gender: 🗌 Male 🔲 Female
Address:			_City, State, ZIP C	Code:	
Preferred Contact Methods: Delow)	Phone (to primary # prov	vided below) [Text (to cell # p	rovided below)) 🗌 Email (to email provided
Note: Carrier charges may apply	. By providing the phone	number(s) and	l email address al	bove, you are c	onsenting to receive
automated calls, emails and/or t	ext messages from CVS	Specialty® abo	ut your prescription	on(s), account,	and health care. Standard data
rates apply. Message frequency					
Primary Phone:			Alternate Phone	:	
					nguage:
Parent/Caregiver/Legal Guardia	ın Name (Last, First):		Relationship t	io patient:	
2 PRESCRIBER INFORM	IATION				
Prescriber's Name: NPI #: DEA #:			State License #	#:	
NPI #: DEA #:	Group or Hos	spital:			
Address:Phone:		City,	State, ZIP Code: _		
Phone:	FaxC	contact Person	:	Contact's P	hone:
3 INSURANCE INFORMA	FION Please fax copy of p	prescription and	d insurance cards	with this form, if	favailable (front and back)
	CAL INFORMATION				
4 DIAGNOSIS AND CLINI Needs by Date:	Ship to: Patient	■] Office [Oth	er:		
Diagnosis (ICD-10):					
K50.00 Crohn's Disease of S	_				
K51.90 Ulcerative colitis, uns		•			
L40.50 Arthropathic Psoriasi	-				
L40.54 Juvenile Psoriatic Art					
M06.9 Rheumatoid Arthritis,					
M08.00 Juvenile Idiopathic A	•				
M32.1 Systemic lupus erythe					
M32.14 Glomerular disease in		natosus			
M45.9 Ankylosing Spondyliti					
M45.A0 Non-Radiographic A		-			
Other Code:					
Patient Clinical Information	<u> </u>				
		IKDA W	/eight: 🔲 k	kg 🔲 lb Height:	
Allergies: New to ther	apy Continuati	on of therapy; [ate of last treatme	ent//	
TB Test Date//_ Dos	sitive Negative		is status:		
Prior therapy, treatment dates, an	d reason(s) for discontinua	ation:			
Nursing and Administration	<u>ı:</u>				
First dose administration of mor	oclonal antibodies (mAB	s) should be a	dministered in a c	ontrolled settin	ng (may vary depending upon
medication specific policy).					
For Remicade/Remicade Biosi					
Specialty pharmacy to coordina					_
Site of Care: Home Infusion					
*Home Infusion/Coram AIS: Dile		•	•	nistration/thera	apy teach train.
**Prescriber's Office/Other Infus	sion Clinic: Drug only for	facility admini	stration		

<u> </u>		Please Complete Patient and F		
Patient Name: _		Patient DOB:	Patient Phone:	
Patient Clinical		_		
Allergies:		NKDA W	eight: 🗌 kg 🗌 lb Height: 🔲 c	m 🗌 in
		Continuation of therapy; D		
			s status:	
Prior therapy, tre	atment dates, and re	eason(s) for discontinuation:		
	ON INFORMATION			
MEDICATION	STRENGTH	DOSE &	DIRECTIONS	QUANTITY/REFILLS
_	☐ 80 mg/4 mL	☐ Induction Dose: Infuse 4 mg/kg eve	erv 4 weeks	Quantity:
Actemra Actemra	200 mg/10 mL	Maintenance Dose: Infuse 8 mg/kg		Quantity: Refills:
	☐ 400 mg/20 mL		Keillis.	
		Ankylosing Spondylitis Induction Do	ose: Infuse IV at 5 mg/kg	
		(Dose =mg) at weeks 0, 2, 6 and e	very 6 weeks thereafter	
		Ankylosing Spondylitis Maintenance	<u>e Dose</u> : Infuse IV at 5 mg/kg	
		(Dose =mg) every 6 weeks		
			c ≥ 6 years old) <u>Induction Dose</u> : Infuse IV at	
		5 mg/kg (Dose =mg) at weeks	-	
		Crohn's Disease (Adult) Maintenance (Dose = mg) every 8 weeks	ce Dose: Infuse IV at 5-10 mg/kg	
		Crohn's Disease (Pediatric ≥ 6 years	s old) Maintenance Dose:	
		Infuse IV at 5 mg/kg (Dose =n	·	
		Plaque Psoriasis & Psoriatic Arthritis		Quantity:
☐ Avsola	100 mg vial	(Dose =mg) at weeks 0, 2, 6 ar		# of 100 mg vial(s)
		Plaque Psoriasis & Psoriatic Arthritis		Refills:
		Infuse IV at 5 mg/kg (Dose =n		
		Rheumatoid Arthritis Induction Dose		
		(Dose =mg) at weeks 0, 2, 6 and		
		Rheumatoid Arthritis Maintenance I		
		(Dose =mg) every 4, 6 or 8 we		
		Ulcerative Colitis (Adult and Pediatr 5 mg/kg (Dose =mg) at weeks		
		Ulcerative Colitis (Adult and Pediatr		
		at 5 mg/kg (Dose =mg) every		
	☐ 120 mg 5 mL	at a ring, kg (Base =ring, avar)	o weeks	
□ p t t.	vial	☐ Induction Dose: 10 mg/kg IV (Dose	=mg) at 2-week intervals for the first 3	Quantity: vials
Benlysta	☐ 400 mg 20 mL	doses and at 4-week intervals thereafte	Refills:	
vial	vial			
	125 mg/5 mL vial	Loading Dose:		
				Quantity:
		Infuse 6 mg/kg (Dose = mg) at Week 0		Refills: 0
☐ Cosentyx		Maintenance Dose: mg) every 4 weeks (max. maintenance dose:		
				Quantity:
				Refills:
	000	300 mg per infusion)		0
300 mg in a single		☐ Induction Dose: 300 mg infused IV over 30 minutes at 0, 2 and 6 weeks, then every 8 weeks thereafter		Quantity: Refills:
— ,	dose vial in individual carton	Maintenance Dose: 300 mg infused IV over 30 minutes every 8 weeks		Remis.
	Strength:		<u> </u>	Quantity:
Other	Suerigui.	Dose:		Refills:
PRESCRIP	ER SIGNATI IPF	REQUIRED (STAMP SIGNAT	URE NOT ALI OWED)	
		•	,	
"Dispense As Writte DAW / May Not Sub	•	essary / Do Not Substitute / No Substitution /	May Substitute / Product Selection Permitted / Substitution Permissible	
•	ignature:	Date:	Prescriber's Signature:	Date:
	.gw.w. V			

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing above, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

Potiont Name:		Please Complete Patient and Prescriber Information	
	 ie:	Patient DOB: Patient Phone: Prescriber Phone:	
Prescriber Nam Patient Clinica		Prescriber Phone:	
Allergies: Treatment statu TB Test Date Prior therapy, tre	s: New to therap	reason(s) for discontinuation:	
	ON INFORMATION		
MEDICATION	STRENGTH	DOSE & DIRECTIONS	QUANTITY/REFILLS
☐ Inflectra	100 mg vial	Ankylosing Spondylitis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 6 weeks thereafter Ankylosing Spondylitis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 6 weeks Crohn's Disease (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Crohn's Disease (Adult) Maintenance Dose: Infuse IV at 5-10 mg/kg (Dose =mg) every 8 weeks Crohn's Disease (Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks Plaque Psoriasis & Psoriatic Arthritis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Plaque Psoriasis & Psoriatic Arthritis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks Rheumatoid Arthritis Induction Dose: Infuse IV at 3 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Rheumatoid Arthritis Maintenance Dose: Infuse IV at 3-10 mg/kg (Dose =mg) every 4, 6 or 8 weeks (circle one) Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks	Quantity: # of 100 mg vial(s) Refills:
Omvoh	300 mg/15 mL single dose vial	Induction Dose ☐ Week 0: Infuse 300 mg via IV infusion over at least 30 minutes ☐ Week 4: Infuse 300 mg via IV infusion over at least 30 minutes ☐ Week 8: Infuse 300 mg via IV infusion over at least 30 minutes	Quantity: Refills: 0 1 Vial 2 Vials 3 Vials
☐ Orencia ☐ Remicade ☐ Renflexis	250 mg vial	☐ Infuse mg at weeks 0, 2 and 4, then every 4 weeks thereafter ☐ Ankylosing Spondylitis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 6 weeks thereafter ☐ Ankylosing Spondylitis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 6 weeks ☐ Crohn's Disease (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter ☐ Crohn's Disease (Adult) Maintenance Dose: Infuse IV at 5-10 mg/kg (Dose =mg) every 8 weeks ☐ Crohn's Disease (Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks ☐ Plaque Psoriasis & Psoriatic Arthritis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter ☐ Plaque Psoriasis & Psoriatic Arthritis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks ☐ Rheumatoid Arthritis Induction Dose: Infuse IV at 3 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter ☐ Rheumatoid Arthritis Maintenance Dose: Infuse IV at 3-10 mg/kg (Dose =mg) every 4, 6 or 8 weeks (circle one) ☐ Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter ☐ Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks	Quantity: Refills: Quantity: # of 100 mg vial(s) Refills:
Other	Strength:	□ Dose:	Quantity: Refills:
"Dispense As Writt DAW / May Not Su Prescriber's S	en" / Brand Medically Ned bstitute ignature:	Date:ATTN: New York and lowa providers, ples	

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing above, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

	_	Please Complete Patient and F	Prescriber Information	
Patient Name: _			Patient Phone:	
Prescriber Name			escriber Phone:	
Patient Clinical	Information:			
Allergies:		NKDA W	'eight: 🗌 kg 🗌 lb Height:	☐ cm ☐ in
		Continuation of therapy; D		
			s status:	
	ON INFORMATIO			
MEDICATION	STRENGTH	DOSE &	DIRECTIONS	QUANTITY/REFILLS
☐ Riabni ☐ Rituxan ☐ Ruxience	☐ 100 mg/10 mL vial ☐ 500 mg/50 mL vial	☐ Infuse two doses of 1000 mg separ	rated by 2 weeks	Quantity: Refills:
Saphnelo	300 mg/2 mL (150 mg/mL)	300 mg IV over a 30-minute period	d, every 4 weeks	Quantity: vials Refills:
Simponi ARIA 50 mg/4 mL single dose vial	Adult RA, PsA, AS Induction Dose Week 0: Infuse 2 mg/kg IV (Dose= Week 4: Infuse 2 mg/kg IV (Dose=	=	Quantity: vials Refills:0 Quantity: vials Refills:0	
	-	Adult RA, PsA, AS Maintenance Dose Infuse 2 mg/kg IV (Dose=mg) over 30 minutes every 8 weeks		Quantity: vials Refills:
		Peds JIA or PsA (≥2 years old) Induction Dose ☐ Week 0: Infuse 80 mg/m² IV (Dose=mg) over 30 minutes ☐ Week 4: Infuse 80 mg/m² IV (Dose=mg) over 30 minutes		Quantity: vials Refills:O Quantity: vials Refills:O
		Peds JIA or PsA (>2 years old) Maint ☐ Infuse 80 mg/m² IV (Dose=m		Quantity: vials Refills:
Skyrizi	600 mg/10 mL (60 mg/mL) single dose vial	Induction Dose: Week 0: Infuse 600 mg IV over at least one hour Week 4: Infuse 600 mg IV over at least one hour Week 8: Infuse 600 mg IV over at least one hour		Quantity: 2 Vials 3 Vials 4 Vials Refills: 0
Stelara	130 mg/26 mL (5 mg/mL) IV single-dose vial	Single IV Induction Dose: 55 kg or less 260 mg at week 0: # of vials to be used 2 more than 55 kg to 85 kg 390 mg at week 0: # of vials to be used 3 more than 85 kg 520 mg at week 0: # of vials to be used 4		Quantity: 2 Vials 3 Vials 4 Vials Refills: 0
☐ Truxima	☐ 100 mg/10 mL vial ☐ 500 mg/50 mL vial	☐ Infuse two doses of 1000 mg separated by 2 weeks		Quantity: Refills:
Other	Strength:	Dose:		Quantity: Refills:
	_	REQUIRED (STAMP SIGNAT	LIDE NOT ALLOWED	
"Dispense As Writte DAW / May Not Sub Prescriber's Si	en" / Brand Medically Necessostitute ignature:	eary / Do Not Substitute / No Substitution /	May Substitute / Product Selection Permitted / Substitution Permissible Prescriber's Signature: ATTN: New York and Iowa providers,	Date:

L. The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing above, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

Autoimmune IV Enrollment Form Nursing Orders

ationt Nomes:			Prescriber Information	
atient Name:		Patient DOB:	Patient Phone: Prescriber Phone:	
escriber Name:	•	P	rescriber Phone:	
atient Clinical Informat			Veight: 🗌 kg 🔲 lb Height:	□om□in
eatment status: \(\int \text{New} \)	to therapy	Continuation of therapy: I	Veight:	
Test Date//	Positive N	egative Henati	tis status:	
) for discontinuation:		
PRESCRIPTION INFO			VILL ONLY BE SENT FOR INFUSIONS DON	IE AT HOME/CODAM AIS**
MEDICATION/SUPPLIES			ENGTH/ DIRECTIONS	QUANTITY/REFILLS
ILDICATION/ SOPPLIES	ROOTE		on drug admin days – SASH or PRN to	QUANTIT I/REFIELS
Catheter: PIV PORT CVC/PICC	IV	maintain IV access and pate PIV: NS 5 mL (Heparin 10 uni CVC/PICC: NS 10 mL & H 3-5 mL.	ncy its/mL 3-5 mL if multiple days) Ieparin 10 units/mL or 100 units/mL access PORT w/ huber needle	Quantity: Refills:
		THE TO THE ATTORACTION	oo ama me o ome.	Hydration max infusion
Hydration: □ NS □ D5W	IV	Pre: 500 mL 1000 mL Concurrent: 500 mL 1000 mL Post: 500 mL 1000 mL	000 mL Other:	rate mL/hr (Adult max rate 250 mL/hr unless otherwise indicated)
☐Epinephrine *nursing requires**	□ IM □ SC	1:1000, 0.3mg/0.3 mL (gr 1:1000, 0.15mg/0.3 mL (1 1:1000, 0.1 mg/kg, Max 0 Mild-Moderate Reactions. M for severe allergic reaction, a	5-30 kg/33-66 lbs) .3mg (under 15kg) 1ay repeat in 3-5 minutes as needed	Quantity: Refills:
Diphenhydramine Oral	РО	Premedication: 12.5 mg/kg (0-30 kg) 25 mg 50 mg (Over 30 kg)		Quantity: Refills:
Diphenhydramine one one one one one one one one one o	Slow IV	1 mg/kg (under 15 kg) 12.5 mg-50 mg (15-30 kg) 25 mg-50 mg (Over 30 kg	g) nay repeat in 3-5 minutes as needed ay)	Quantity: Refills:
☐ Flush Orders:	Peripheral Access Central Venous Access	☐ 10 mL NS post flush ☐ 50 mL NS post flush to clear medication from tubing (recommended if no post-hydration) ☐ Other:		Send quantity sufficient for medication days supply
Additional				
ledication:				
Patient is interested in patient supp		STAMP SIGNATURE NOT ALLOWED JIRED (STAMP SIGNAT	, ,,	provided as needed for administration
"Dispense As Written" / Brand M DAW / May Not Substitute	edically Necessary / D	o Not Substitute / No Substitution /	May Substitute / Product Selection Permitted / Substitution Permissible Prescriber's Signature:	Date:
Prescriber's Signature:				

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing above, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA

request as my signature.

CONFIDENTIALITY NOTICE: This communication and any attachments may contain confidential and/or privileged information for the use of the designated recipients named above. If you are not the intended recipient, you are hereby notified that you have received this communication in error and that any review, disclosure, dissemination, distribution or copying of it or its contents is prohibited. If you have received this communication in error, please notify the sender immediately by telephone and destroy all copies of this communication and any attachments.

Plan member privacy is important to us. Our employees are trained regarding the appropriate way to handle members' private health information.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Specialty and/or one of its

affiliates.